

Graph-truncations of 3-polytopes.

Nickolai Erokhovets *

Abstract

In this paper we study the operation of cutting off edges of a simple 3-polytope P along the graph Γ . We give the criterion when the resulting polytope is simple and when it is flag. As a corollary we prove the analog of Eberhard's theorem about the realization of polygon vectors of simple 3-polytopes for flag polytopes.

1 Introduction.

For the introduction to the polytope theory we recommend the books [Gb03, Z07].

Definition 1.1. A *convex polytope* P is a set

$$P = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}_i \mathbf{x} + b_i \geq 0, i = 1, \dots, m\}$$

Let this representation be *irredundant*, that is deletion of any inequality changes the set. Then each hyperplane $\mathcal{H}_i = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{a}_i \mathbf{x} + b_i = 0\}$ defines a *facet* $F_i = P \cap \mathcal{H}_i$.

In the following by a *polytope* we mean a convex polytope.

A *dimension* $\dim(P)$ of the polytope P is defined as $\dim \text{aff}(P)$. We will consider n -dimensional polytopes (n -polytopes) in \mathbb{R}^n .

A *face* F of a polytope is an intersection $F = P \cap \{\mathbf{a}\mathbf{x} + b = 0\}$ for some *supporting hyperplane* $\{\mathbf{a}\mathbf{x} + b = 0\}$, i.e. $\mathbf{a}\mathbf{x} + b \geq 0$ for all $\mathbf{x} \in P$. Each face is a convex polytope itself. 0-dimensional faces are called *vertices*, 1-dimensional faces – *edges*, $(n-1)$ -faces – *facets*. It can be shown that the set of all facets is $\{F_1, \dots, F_m\}$. Intersection of any set of faces of polytope is a face again (perhaps empty).

A vertex of an n -polytope P is called *simple* if it is contained in exactly n facets. An n -polytope P is called *simple*, if all its vertices are simple. Each k -face of a simple polytope is an intersection of exactly $n-k$ facets.

A *combinatorial polytope* is an equivalence class of combinatorially equivalent convex polytopes, where two polytopes are *combinatorially equivalent* if there is an inclusion-preserving bijection of the sets of their faces.

A simple polytope is called *flag* if any set of pairwise intersecting facets $F_{i_1}, \dots, F_{i_k} : F_{i_s} \cap F_{i_t} \neq \emptyset$ has nonempty intersection $F_{i_1} \cap \dots \cap F_{i_k} \neq \emptyset$.

A *non-face* is the set $\{F_{i_1}, \dots, F_{i_k}\}$ with $F_{i_1} \cap \dots \cap F_{i_k} = \emptyset$. A *missing face* is an inclusion-minimal non-face.

The following results are well-known.

Proposition 1.2. A polytope P is flag if and only if all its missing faces have cardinality 2.

Proposition 1.3. Each face of a flag polytope is a flag polytope again.

Proposition 1.4. The simplex Δ^n is not flag for $n \geq 3$. A 3-polytope $P^3 \neq \Delta^3$ is not flag if and only if it has missing face of cardinality 3: $\{F_i, F_j, F_k\}$, $F_i \cap F_j, F_j \cap F_k, F_k \cap F_i \neq \emptyset$, $F_i \cap F_j \cap F_k = \emptyset$.

Definition 1.5. Missing face $\{F_i, F_j, F_k\}$ of a 3-polytope P^3 we will also call a 3-belt.

Let $f_i(P)$ be the number of i -faces of the polytope P .

*The work was supported by RFBR grant No 14-01-31398-a

Proposition 1.6 (The Euler formula). *For a 3-polytope we have*

$$f_0 - f_1 + f_2 = 2$$

Let p_k be the number of 2-faces of P that are k -gons.

Proposition 1.7. *For a simple 3-polytope P^3 we have*

$$3p_3 + 2p_4 + p_5 = 12 + \sum_{k \geq 7} (k-6)p_k \quad (*)$$

Proof. Let us count the number of pairs (edge, it's vertex). It is equal to $2f_1$, and since P is simple to $3f_0$. Then $f_0 = \frac{2f_1}{3}$ and from the Euler formula we obtain $2f_1 = 6f_2 - 12$. Then counting the pairs (facet, it's edge) we have

$$\sum_{k \geq 3} kp_k = 2f_1 = 6 \left(\sum_{k \geq 3} p_k \right) - 12,$$

which implies the formula (*). \square

Theorem 1.8 (Eberhard). [Eb1891]

For every sequence $(p_k | 3 \leq k \neq 6)$ of nonnegative integers satisfying (), there exist values of p_6 such that there is a simple 3-polytope P^3 with $p_k = p_k(P^3)$ for all $k \geq 3$.*

2 Graph-truncations

Construction 2.1. Consider a subgraph Γ without isolated vertices in the edge-vertex graph $G(P)$ of a simple 3-polytope P . For each edge $E_{i,j} = F_i \cap F_j = P \cap \{\mathbf{x} \in \mathbb{R}^3 : (\mathbf{a}_i + \mathbf{a}_j)\mathbf{x} + (b_i + b_j) = 0\}$ consider the halfspace $\mathcal{H}_{i,j,\varepsilon}^+ = \{\mathbf{x} \in \mathbb{R}^3 : (\mathbf{a}_i + \mathbf{a}_j)\mathbf{x} + (b_i + b_j) \geq \varepsilon\}$. Set

$$P_{\Gamma,\varepsilon} = P \cap \bigcap_{E_{i,j} \in \Gamma} \mathcal{H}_{i,j,\varepsilon}^+$$

For small values of ε the combinatorial type of $P_{\Gamma,\varepsilon}$ does not depend on ε . We will denote it P_Γ and call a *graph-truncation* of P .

Facets of the polytope P_Γ are in one-to one correspondence with facets F_i of P (denote such facets by the same symbol F_i) and edges $F_i \cap F_j \in \Gamma$ (denote such facets as $F_{i,j}$).

Proposition 2.2. *The polytope P_Γ is simple if and only if the graph Γ does not contain vertices of valency 2.*

Proof. For small ε all new vertices of the polytope $P_{\Gamma,\varepsilon}$ lie in small neighborhoods of vertices of P . Consider a vertex $\mathbf{v} = F_{i_1} \cap F_{i_2} \cap F_{i_3}$ of P and introduce new coordinates in \mathbb{R}^3 by the formulas

$$y_1 = \mathbf{a}_{i_1}\mathbf{x} + b_{i_1}, \quad y_2 = \mathbf{a}_{i_2}\mathbf{x} + b_{i_2}, \quad y_3 = \mathbf{a}_{i_3}\mathbf{x} + b_{i_3}.$$

In new coordinates in some neighborhood $U(\mathbf{v})$ of $\mathbf{v} = \mathbf{0}$ the polytope $P_{\Gamma,\varepsilon}$ has irredundant representation

$$P_{\Gamma,\varepsilon} \cap U(\mathbf{v}) = \{\mathbf{y} \in \mathbb{R}^3 : y_1, y_2, y_3 \geq 0; y_p + y_q \geq \varepsilon, \text{ if } F_{i_p} \cap F_{i_q} \in \Gamma\}$$

The polytope $P_{\Gamma,\varepsilon}$ has non-simple vertex in $U(\mathbf{v})$ if and only if there is some point in $P_{\Gamma,\varepsilon} \cap U(\mathbf{v})$ that belongs to 4 facets.

If $\mathbf{v} \notin \Gamma$ then $\mathbf{v} \in P_{\Gamma,\varepsilon}$ is the only vertex in $U(\mathbf{v})$ and it is simple.

If \mathbf{v} has valency 1 in Γ , say $F_{i_1} \cap F_{i_2} \in \Gamma$, and some point lies in $F_{i_1}, F_{i_2}, F_{i_3}, F_{i_1,i_2}$, then $y_1 = y_2 = y_3 = 0$ and $y_1 + y_2 = \varepsilon$. Contradiction.

If \mathbf{v} has valency 2 in Γ , say $F_{i_1} \cap F_{i_2}, F_{i_2} \cap F_{i_3} \in \Gamma$, then the point $(0, \varepsilon, 0) \in P_{\Gamma,\varepsilon}$ belongs to $F_{i_1}, F_{i_3}, F_{i_1,i_2}, F_{i_2,i_3}$, so $P_{\Gamma,\varepsilon}$ is not simple.

Let \mathbf{v} has valency 3 in Γ and some point belongs to 4 facets. If there are F_{i_1,i_2}, F_{i_2,i_3} and F_{i_3,i_1} , among them, then $y_1 = y_2 = y_3 = \frac{\varepsilon}{2}$, and there can not be neither F_{i_1} , nor F_{i_2} , nor F_{i_3} . Therefore there should be at least two of $F_{i_1}, F_{i_2}, F_{i_3}$, say F_{i_1}, F_{i_2} . Then $y_1 = y_2 = 0$. But $y_1 + y_2 \geq \varepsilon$. Contradiction. So $P_{\Gamma,\varepsilon}$ has only simple vertices in $U(\mathbf{v})$ in this case. \square

Theorem 2.3. *A simple 3-polytope P_Γ is flag if and only if any triangular facet of P contains no more than one edge in Γ and for any 3-belt (F_i, F_j, F_k) of P one of the edges $F_i \cap F_j$, $F_j \cap F_k$, $F_k \cap F_i$ belongs to Γ .*

Proof. Since P_Γ is simple, Proposition 2.2 implies that valency of each vertex of Γ is 1 or 3.

If P contains a 3-belt (F_i, F_j, F_k) , such that $F_i \cap F_j$, $F_j \cap F_k$, $F_k \cap F_i \notin \Gamma$, then (F_i, F_j, F_k) is either a 3-belt in $P_{\Gamma, \varepsilon}$. Consider a triangular face of P . If exactly two of its edges belong to Γ , then Proposition 2.2 implies that valency of their common vertex is 3 and other vertices have valency 1 in Γ . If all three edges belong to Γ , then all their vertices have valency 3 in Γ . In both cases after truncation the face remains to be triangular, so $P_{\Gamma, \varepsilon}$ is not flag. Thus we proved the only if part of the theorem.

$P_{\Gamma, \varepsilon} \neq \Delta^3$, since it contains more than 4 facets. Therefore if it is not flag, then there is a 3-belt (G_1, G_2, G_3) in $P_{\Gamma, \varepsilon}$ by Proposition 1.4.

If $G_1 = F_i$, $G_2 = F_j$, $G_3 = F_k$, then either (F_i, F_j, F_k) is a 3-belt in P , or there is a vertex $v = F_i \cap F_j \cap F_k \in P$. In the first case one of the edges $F_i \cap F_j$, $F_j \cap F_k$, or $F_k \cap F_i$ belongs to Γ and is cut off when we pass to $P_{\Gamma, \varepsilon}$, so the corresponding facets do not intersect in $P_{\Gamma, \varepsilon}$. In the second case the vertex v is cut off, since $F_i \cap F_j \cap F_k = \emptyset$ in $P_{\Gamma, \varepsilon}$. It is possible only if we cut off one of the edges containing this vertex, so the corresponding two facets do not intersect in $P_{\Gamma, \varepsilon}$.

If $G_1 = F_i$, $G_2 = F_j$, and $G_3 = F_{p,q}$ correspond to an edge $E_{p,q} = F_p \cap F_q$ of P , then $F_i \cap F_j \neq \emptyset$ and $E_{p,q}$ intersects both F_i and F_j . Since $F_i \cap F_j$ was not cut off, we have $\{i, j\} \neq \{p, q\}$. If $i \in \{p, q\}$ or $j \in \{p, q\}$, then the edge $F_p \cap F_q$ intersects the edge $F_i \cap F_j$ at the vertex, so the facets F_i , F_j , and $F_{p,q}$ have common vertex in $P_{\Gamma, \varepsilon}$. Now let $\{i, j\} \cap \{p, q\} = \emptyset$. Then (F_i, F_j, F_p) or (F_i, F_j, F_q) is a 3-belt in P . Otherwise $F_i \cap F_j \cap F_p \neq \emptyset$, $F_i \cap F_j \cap F_q \neq \emptyset$, $F_p \cap F_q \cap F_i \neq \emptyset$, and $F_p \cap F_q \cap F_j \neq \emptyset$, therefore $P = \Delta^3$, all its facets are triangles and in any triangle no more than one edge is cut off. Since $F_i \cap F_j \notin \Gamma$ and $F_p \cap F_q \in \Gamma$, in facets F_p and F_q the only edge $F_p \cap F_q$ is cut off and Γ contains no other edges. Then the facets F_p and F_q are triangles in $P_{\Gamma, \varepsilon}$ either, and it is not flag. By assumption one of the edges of the 3-belt we obtain belongs to Γ . Since the edge $F_i \cap F_j$ was not cut off, one of the edges $F_i \cap F_p$, $F_j \cap F_p$, $F_i \cap F_q$ and $F_j \cap F_q$ belongs to Γ and was cut off, say $F_i \cap F_p$. Then $F_i \cap F_{p,q} = \emptyset$, which is a contradiction.

If only one of the facets (G_1, G_2, G_3) corresponds to a facet of P , say $G_1 = F_i$, then two other facets correspond to edges of P that both intersect F_i and have common vertex. If both edges belong to F_i , then $F_i \cap G_2 \cap G_3 \neq \emptyset$. If exactly one of them belongs to F_i , say corresponding to G_2 , then $F_i \cap G_3 = \emptyset$. At last, if both of them do not belong to F_i , then their common vertex v does not belong to F_i , and these two edges and their common vertex define some facet F_j that has with F_i two common vertices – the remaining ends of two edges, thus $F_i \cap F_j$ is an edge, connecting these vertices. Then F_j is a triangle containing two edges in Γ . Contradiction.

At last if all three facets of 3-belt correspond to edges of P , then these edges pairwise intersect. Two of them define some facet F_i . If the third edge does not belong to F_i , then all three edges have common vertex and in $P_{\Gamma, \varepsilon}$ the corresponding facets have a common vertex either. If the third edge belongs to F_i , then F_i is a triangle with three edges in Γ . Contradiction.

Thus we have considered all possible cases, and the theorem is proved. \square

3 Application

As an application of Theorem 2.3 we prove an analog of Eberhard's theorem for flag 3-polytopes. Since any face of a flag polytope is flag itself, we have $p_3(P^3) = 0$ for any flag polytope.

Theorem 3.1. *For every sequence $(p_k | 3 \leq k \neq 6)$ of nonnegative integers satisfying $p_3 = 0$ and (*), there exist values of p_6 such that there is a flag simple 3-polytope P^3 with $p_k = p_k(P^3)$ for all $k \geq 3$.*

Proof. From Eberhard's Theorem 1.8 it follows that there exist values of p_6 such that there is a polytope P^3 with $p_k = p_k(P)$ for all $k \geq 3$. Let us consider the graph $\Gamma = G(P^3)$. Since $p_3 = 0$, we obtain from Theorem 2.3 that $P_{G(P)}$ is flag. On the other hand, facets of $P_{G(P)}$ are in one-to-one correspondence with facets and edges of P . Moreover, k -gonal facets of P correspond to k -gonal facets of $P_{G(P)}$ and edges of P correspond to 6-gonal facets of $P_{G(P)}$. Therefore

$$p_k(P_{G(P)}) = \begin{cases} p_k(P), & k \neq 6; \\ p_6(P) + f_1(P), & k = 6. \end{cases}$$

This proves the theorem. □

The author is grateful to professor V.M. Buchstaber for encouraging discussions and a permanent attention to his work.

References

- [Eb1891] V. Eberhard, *Zur Morphologie der Polyheder*, Leipzig 1891.
- [Gb03] Branko Grünbaum, *Convex polytopes*, Graduate texts in Mathematics 221, Springer-Verlag, New York, 2003.
- [Z07] Günter M. Ziegler, *Lectures on polytopes*, Graduate texts in Mathematics 152, Springer, 2007.